

A P P L I C A T I O N

for

UNITED STATES LETTERS PATENT

on

A METHOD FOR DETERMINING THE PROGNOSIS OF CANCER PATIENTS
BY MEASURING LEVELS OF BAG EXPRESSION

by

John C. Reed

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"EXPRESS MAIL" MAILING LABEL NUMBER: EL 454 952 867 US

DATE OF DEPOSIT: July 9, 1999

Sheets of Drawings: One
Docket No.: P-LJ 3578

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS BEING
DEPOSITED WITH THE UNITED STATES POSTAL SERVICE
"EXPRESS MAIL POST OFFICE TO ADDRESSEE" SERVICE UNDER
37 C.F.R. 1.10 ON THE DATE INDICATED ABOVE, AND IS
ADDRESSED TO: THE ASSISTANT COMMISSIONER FOR PATENTS
WASHINGTON, D.C. 20231.

Rick de la Torre
(TYPED OR PRINTED NAME OF PERSON MAILING PAPER OR FEE)

Rick de la Torre
(SIGNATURE OF PERSON MAILING PAPER OR FEE)

Attorneys

CAMPBELL & FLORES LLP
4370 La Jolla Village Drive, Suite 700
San Diego, California 92122

**A METHOD FOR DETERMINING THE PROGNOSIS OF CANCER PATIENTS
BY MEASURING LEVELS OF BAG EXPRESSION**

Portions of the invention described herein were made in the course of research supported in part by NIH grant CA-67329. The Government may have certain rights in this invention.

FIELD OF THE INVENTION

The present invention relates generally to determining the risk of progression or spread of cancers.

10

BACKGROUND INFORMATION

Prognosis in clinical cancer is an area of great concern and interest. It is important to know the aggressiveness of the malignant cells and the likelihood of tumor recurrence or spread in order to plan the most effective therapy. Breast cancer, for example, is managed by several alternative strategies. One of every nine women currently develops breast cancer at some point in her life. In some cases local-regional therapy is utilized, consisting of mastectomy or lumpectomy with or without radiation, while in other cases when spread of disease is detected or suspected, systemic therapy is instituted, such as chemotherapy or hormonal therapy. Among women with early-stage breast cancers treated with lumpectomy and local radiotherapy, 10-20% will experience local recurrences and 30-40% will develop distant metastatic disease which is often fatal (Fischer et al. 1991, Lancet 338:327-331).

Current treatment decisions for individual breast cancer patients are frequently based on (1) the number of axillary lymph nodes involved with disease, (2) estrogen receptor and progesterone receptor status, (3) the size of the primary tumor, and (4) stage of disease at diagnosis. It has also been reported that DNA aneuploidy and proliferative rate (percent S-phase) can help in predicting the course of disease. In addition, the overexpression of the HER2/Neu oncoprotein has been shown to predict breast cancer patients at risk for metastatic disease and novel therapeutic strategies have been developed to target this receptor (Slamon et al. U.S. Patent No. 4,968,603; Slamon et al. 1989, Science 244:707-712). However, even with these additional factors, practitioners are still unable to accurately predict the course of disease for all breast cancer patients. There is clearly a need to identify new markers, in order to separate patients with good prognosis who may not require further therapy from those more likely to recur who might benefit from more intensive treatments.

This is particularly true in the case of breast cancer which has not progressed to the axillary lymph nodes. There is now evidence in prospective randomized clinical trials that adjuvant endocrine therapy and adjuvant chemotherapy beginning immediately after surgical removal of the primary breast tumor can be of benefit in some of these node-negative patients. This has led to official and unofficial recommendations that most if not all node-negative breast cancer patients should be considered for some form of adjuvant therapy. But since the majority (about 70%) of these patients enjoy long-term survival following surgery and/or radiotherapy without further treatment, it may be

inappropriate to recommend adjuvant therapy for all of these patients. Accordingly, there is a need for methods to distinguish those node-negative patients who are "cured" from those destined to recur, such that only the latter are treated. Thus, there is a great need for a general method of predicting tumor recurrence or spread in these patients and in cancer patients in general once a primary tumor is detected. The present invention satisfies these needs and provides related advantages as well.

SUMMARY OF THE INVENTION

In accordance with the present invention, there are provided methods for determining a prognosis of disease free or overall survival in a patient suffering from cancer. In one embodiment, it has been found that high levels or "overproduction" of BAG proteins in primary tumor tissue show an unexpected and surprisingly high correlation with lack of tumor recurrence or spread, and therefore long-term disease free or overall survival. Thus, the present invention advantageously provides a significant advancement in cancer management because early identification of patients at risk for tumor recurrence or spread will permit aggressive early treatment with significantly enhanced potential for survival.

Also provided are methods for predicting the risk of tumor recurrence or spread in an individual having a cancer tumor; methods for screening a cancer patient to determine the risk of tumor metastasis; methods for determining the proper course of treatment for a patient suffering from cancer; and kits for use in practicing the invention methods.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows that high BAG-1 Protein levels are associated with longer patient survival.

5 Kaplan-Meier survival curves containing the proportion of early stage breast cancer patients whose tumors contain high (O; referred to in the figure as BAG-1 positive) versus low (●; referred to as BAG-1 negative) levels of BAG-1 protein is plotted against time (years) for overall
10 survival (OS) (top) and distant metastasis-free survival (DMFS) (bottom).

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, methods are provided for determining a prognosis of
15 disease free or overall survival in a patient suffering from cancer, said method comprising:

- (a) determining a BAG gene expression level in a cancerous tissue sample from said patient; and
- 20 (b) classifying said patient as belonging to either a first or second group of patients, wherein said first group of patients having high levels of expression of the BAG gene is classified as having a different likelihood of
25 suffering tumor recurrence or spread than said second group of patients having low levels of expression of the BAG gene.

The invention method takes advantage of a newly discovered correlation between abnormal levels of BAG
30 expression in cancerous cells, and the likelihood of cancer-free survival in patients (see, e.g., the Examples

section). The invention method comprises determining the level of expression of BAG in the cancerous tissue of a patient. The patient may then be classified as belonging to a first group of patients having high levels of
5 expression of BAG, or alternatively, to a second group of patients having low levels of expression of BAG. Determination of the prognosis for the patient suffering from cancer may be made by determining whether the group to which the patient has been assigned correlates with a
10 higher or lower likelihood of disease-free or overall survival with respect to the group to which the patient was not assigned.

For example, in one embodiment of the invention, it has been discovered that overproduction or
15 a high level of expression of a BAG gene correlates to patients having a decreased risk of tumor recurrence or spread (e.g., local recurrence or spread and/or metastasis). Thus, in this embodiment, patients belonging to a first group having high levels of
20 expression of a BAG gene are classified as having a decreased risk of tumor recurrence or spread compared to a second group of patients having low levels of expression of a BAG gene. This embodiment is particularly useful in evaluating the prognosis of breast
25 cancer patients.

For example, in this embodiment, when BAG gene expression exceeds a determined basal level (also referred to herein as a reference level), it has been found to become a significant factor in decreased tumor
30 recurrence or spread. When tumor cell determined basal levels are exceeded, a BAG expression level is characterized as "high" or "overproduced" and indicates, particularly in the case of breast cancer, a decreased

risk of tumor recurrence or spread. Whereas, the mere presence, without overproduction, of BAG protein expression has not previously been correlated with any risk of tumor recurrence or spread.

5 In another embodiment of the invention, high levels of expression of a BAG gene correlate to patients having an increased risk of tumor recurrence or spread. Thus, in this embodiment, patients belonging to a first group of patients having high levels of expression of the
10 BAG gene are classified as having an increased risk of tumor recurrence or spread compared to a second group of patients having low levels of expression of the BAG gene.

 In yet another embodiment of the invention, low levels of expression of a BAG gene correlate to patients
15 having a decreased risk of tumor recurrence or spread. Thus, in this embodiment, patients belonging to a second group of patients having low levels of expression of the BAG gene are classified as having a decreased risk of tumor recurrence or spread compared to a first group of
20 patients having high levels of expression of the BAG gene.

 In another embodiment of the invention, low levels of expression of a BAG gene correlate to patients having an increased risk of tumor recurrence or spread.
25 Thus, in this embodiment, patients belonging to a second group of patients having low levels of expression of the BAG gene are classified as having an increased risk of tumor recurrence or spread compared to a first group of patients having high levels of expression of the BAG
30 gene.

The invention methods are useful in the prognosis of disease-free or overall survival of individuals with neoplastic diseases, including both solid tumors and hematopoietic cancers. Exemplary
5 neoplastic diseases include carcinomas, such as adenocarcinomas and melanomas; and sarcomas, such as various leukemias or lymphomas. Of particular interest are breast cancer, prostate cancer, lung cancer, colon cancer, leukemia, lymphoma, and oral cancer; more
10 particularly breast cancer.

As used herein, the terms "cancer recurrence" and "tumor recurrence," and grammatical variants thereof, refer to further growth of neoplastic or cancerous cells after diagnosis of cancer. Particularly, recurrence may
15 occur when further cancerous cell growth occurs in the cancerous tissue. "Tumor spread," similarly, occurs when the cells of a tumor disseminate into local or distant tissues and organs; therefore tumor spread encompasses tumor metastasis.

20 As used herein, the phrase "disease-free survival," refers to the lack of such tumor recurrence and/or spread and the fate of a patient after diagnosis, with respect to the effects of the cancer on the life-span of the patient. The phrase "overall survival"
25 refers to the fate of the patient after diagnosis, despite the possibility that the cause of death in a patient is not directly due to the effects of the cancer. The phrases, "likelihood of disease-free survival", "risk of recurrence" and variants thereof, refer to the
30 probability of tumor recurrence or spread in a patient subsequent to diagnosis of cancer, wherein the probability is determined according to the process of the invention.

Depending on the nature of the cancer, an appropriate patient sample is obtained. As used herein, the phrase "cancerous tissue sample" refers to any cells
5 obtained from a cancerous tumor. In the case of solid tumors which have not metastasized, a tissue sample from the surgically removed tumor will typically be obtained and prepared for testing by conventional techniques. Alternatively, a body fluid sample, such as a lymph,
10 blood or serum sample, or an exudate fluid sample such as the cancerous organ exudate (e.g., exudate from the breast) may be collected and used as the sample to be analyzed. In the case of leukemias, lymphocytes or leukemic cells will be obtained and appropriately
15 prepared. Similarly, in the case of any metastasized cancer, cells may be drawn from a body fluid such as lymphatic fluid, blood, serum, or a distally infected organ or exudate thereof. While BAG levels will typically be measured within the cancerous cells of a
20 patient, levels of BAG will also be measured in a body fluid sample (e.g., serum) as a result of BAG having been secreted or otherwise released from cells (e.g., by cell rupture).

As set forth herein, the invention method for
25 determining the prognosis of a cancer patient comprises: (a) determining the level of BAG gene expression in cancerous tumor cells of a patient, and (b) classifying the patient as having either a low or high likelihood of tumor recurrence or spread based on the level of BAG
30 gene expression in the tumor sample or body fluid sample (e.g., serum) of the patient relative to the reference level.

Determining levels of BAG gene expression

Once the patient tissue or cell sample is obtained, prognosis of disease free or overall survival includes determination of the levels of BAG gene expression or amplification in a cancerous tumor. Measurement of BAG gene expression is performed quantitatively such that the level of gene expression can be determined. The BAG expression level is then used to determine the prognosis of disease-free or overall survival of a cancer patient based on the correlations provided herein. This is possible because the likelihood of tumor recurrence or spread correlates with the level of BAG expression in tumor cells. For example, it has been found that when the level of BAG expression is high, the likelihood of metastasis of certain cancers, such as breast cancer, is low. The level of BAG expression may be used as the sole factor in assessing the disease status, or along with the additional factors, including, in the illustrative case of breast cancer, lymph node status, estrogen receptor status, and the like.

As used herein, "BAG gene" refers to the nucleic acid sequence encoding a protein comprising a BAG domain as set forth in Takayama et al. 1999, Journal of Biological Chemistry 274:781-786, which is incorporated herein in its entirety. Exemplary BAG proteins include the various isoforms, preferably human, of BAG-1 (including isoforms such as BAG-1N, BAG-1M, and BAG-1L, see discussion below), BAG-2, BAG-3, BAG-4, BAG-5 proteins, and any other protein which contains a BAG domain and any mutation or partial deletion thereof, including a mutation or partial deletion which results in enhanced or decreased BAG gene expression or in the expression of overactive, underactive, or inactive BAG

protein. Other BAG proteins include BAG-1 and BAG-2 proteins from *Caenorhabditis elegans*, and BAG-1A and BAG-1B from *Schizosaccharomyces pombe* (Takayama et al. 1999, supra). Thus, as used herein a BAG protein has a
5 conserved 45-amino acid region near the C-terminus, referred to herein as the BAG domain.

A preferred BAG protein for use herein is human BAG-1. BAG-1 protein was originally identified as a novel regulator of apoptosis by virtue of its ability to
10 bind Bcl-2, a potent blocker of cell death. Since its initial characterization, BAG-1 has been found to interact with not only Bcl-2, but also a number of other proteins which are known to be critical in controlling cell death, including Hsp70/Hsc70, Raf-1, HGF receptor,
15 PDGF receptor, and several steroid hormone receptors.

Previous studies have shown that in both mouse and humans, there is only a single size of BAG-1 mRNA molecules transcribed, but a plurality of sizes of BAG-1 polypeptides are translated (Takayama et al. 1998, Cancer
20 Research 58:3116-3131). In mouse, two isoforms have been identified: the normal length BAG-1 (referred to herein as "BAG-1N"); and a longer BAG-1 polypeptide, or BAG-1L. In human, in addition to the presence of the corresponding BAG-1N and BAG-1L isoforms, a medium length
25 BAG-1 polypeptide, referred to as BAG-1M, is also translated. The cDNA complementary to the BAG-1 mRNA for mouse and human are set forth in U.S. Patent 5,539,094 and in Takayama et al. 1998, supra, (each of which are incorporated herein by reference in their entirety). The
30 polypeptide sequences for the two mouse isoforms BAG-1N and BAG-1L are 219 and 355 amino acids long, respectively (see Takayama et al. 1998, supra). The polypeptide

sequences for human BAG-1N, BAG-1M, and BAG-1L are 230, 274 and 345 amino acids in length, respectively.

BAG polypeptides, and particularly BAG-1 (especially BAG-1N and BAG-1L), are present in numerous normal human tissues. Both BAG-1N and BAG-1L have been found to be present at abnormal levels in numerous cancers (Takayama et al., 1998, supra; and Yang et al., 1999, Experimental Cell Research 247:200-207), including for example: colon cancer, leukemia, lymphoma, breast cancer, prostate cancer, lung cancer, melanoma, ovarian cancer, cervical cancer, and renal cancer. However, a correlation of BAG expression levels with prognosis of disease-free or overall survival has heretofore been unavailable.

Determination of BAG gene expression or amplification levels may be performed by one or more of the methods known to one of ordinary skill in the art. For example, BAG amplification or expression levels may be determined by detection of (a) a BAG polypeptide, (b) mRNA encoding a BAG protein, (c) a portion of DNA which constitutes a BAG gene, or (d) any combination thereof.

For example, levels of BAG gene expression can be detected by measuring levels of BAG protein using agents that bind specifically to a BAG polypeptide. As used herein, the term "agent that binds BAG protein" refers to any molecule that specifically binds to a BAG protein, including, for example, BAG-1N, BAG-1L, BAG-1M, and/or polypeptide fragments thereof, and thereby detects the level of BAG expression. Such agents are preferably labeled for detection using methods well known to those of skilled in the art. A variety of agents are contemplated for use herein to specifically detect BAG

protein, preferably BAG-1, including proteins known to bind specifically to BAG, antibodies to BAG (such as described in United States Patent 5,641,866, incorporated herein by reference in its entirety), or peptides which
5 specifically bind BAG. A preferred agent for binding to BAG-1, such as the anti-BAG-1 antibody used herein (See, for example, Takayama et al. 1998, supra, for description of antibody preparation) binds to all isoforms of BAG-1.

Other, non-antibody proteins, may also be used
10 as "agents." For example, BAG proteins are known to specifically bind numerous proteins, such as Bcl-1, Raf-1, HGF-receptor, PDGF-receptor, Hsp70, Hsc70, steroid hormone receptors, and the like. As a result, any of these proteins, or active BAG binding fragments thereof,
15 may be used to specifically bind BAG. An exemplary active binding fragment of a protein which binds BAG-1 (and also BAG-2 and BAG-3) is a BAG binding domain of Hsp70. The ATPase domain of Hsp70 may be expressed in a truncated form, lacking the carboxy-terminal peptide-
20 binding domain. In this form, Hsp70 will not indiscriminantly bind proteins in non-native conformations; however, the ATPase domain of Hsp70 is still capable of binding BAG-1 (or BAG-2 or BAG-3) protein. Therefore, an actively binding fragment of a
25 protein known to specifically bind BAG may be used as an "agent" which specifically binds BAG protein.

Antibodies, both monoclonal and polyclonal, may be used as specifically binding agents which bind BAG protein or a polypeptide fragment thereof. Also
30 contemplated herein as BAG binding agents are any mutants of proteins which specifically bind BAG, whether by deletion (as above exemplified), addition (e.g., addition of a GST domain or a GFP domain), or sequence

modification (e.g., site-specific mutagenesis), and the like.

It is also contemplated herein that one or more of the BAG specific binding agents may be used in a single assay to measure BAG protein levels. For example, a certain protein known to interact with a specific portion of the BAG-1 protein, such as, for example, a first antibody raised against the N-terminal 60 amino acids of the BAG-1L protein, may be coupled with another specifically binding protein, such as a second antibody raised against a sequence contained in both BAG-1N and BAG-1L. Using these two antibodies in a single assay, the specific levels of the differently translated BAG-1 polypeptides may be measured by differentially measuring the two antibodies.

Preparation of the agent for use in the detection of BAG protein levels will be carried out using the methods of one of ordinary skill in the art, such as the methods exemplified in the Current Protocols in Molecular Biology, and in U.S. Patent No. 5,882,864. Similarly, detection of BAG protein levels may be carried out using any of the methods known to one of ordinary skill in the art including histochemical staining, Western Blot analysis, immunoprecipitation (or the equivalent thereof for non-antibody agents), and the like. In a preferred embodiment of the invention, the method of detecting BAG protein levels is an immunoassay (such as an ELISA, immuno-PCR, and the like), which includes the use of at least one antibody. Measurement of the polypeptide encoded by a BAG gene may include measurements of fragments of the polypeptide, wherein the fragments arise from transcriptional or translational variants of the gene; or alternatively, differently sized

polypeptides arise as a result of post translational modifications including proteolysis of a larger portion of a BAG polypeptide.

An exemplary immuno assay for use in the invention methods for detecting BAG protein levels is an immuno-polymerase chain reaction immuno-PCR assay (described in U.S. Patent 5,665,539, which is incorporated herein in its entirety). Immuno-PCR utilizes an antibody (or other agent which binds BAG) to detect the BAG protein, wherein the antibody (or other agent) is linked to a molecule (typically biotin) which specifically binds a bridging molecule (typically avidin), wherein this bridging molecule is capable of binding a second molecule (typically biotin) attached to a nucleic acid marker. This nucleic acid marker is then amplified using PCR methods. This sensitive detection method is particularly useful when BAG levels are often difficult to detect by other methods, for example, detection of BAG in serum.

Measurement of the polypeptide encoded by a BAG gene may further be carried out to specifically measure: (a) the level of BAG produced in the entire cell, (b) the level of BAG produced in the cytosol, (c) the level of BAG produced in the nucleus, (d) level of BAG present in cell-free extract (e.g., serum), and (e) any combination thereof. Exemplary methods which can be used in such measurements include *in situ* methods such as histochemical staining, particularly differential staining between the cytosol and the nucleus, and *in vitro* methods such as Western blot analysis of nuclear extracts, cytosolic extracts, or serum.

Detection of levels of mRNA encoding BAG may also serve as an indicator of BAG expression. The methods used to detect mRNA levels will include the detection of hybridization or amplification with the mRNA
5 encoding BAG. This detection may be carried out by analysis of mRNA either *in vitro* or *in situ* (e.g., in a tissue sample) using one of the methods known to one of ordinary skill in the art as exemplified in the Current Protocols in Molecular Biology (John Wiley & Sons, 1999);
10 in U.S. Patent No. 5,882,864; and the like. A BAG mRNA detected will be any RNA transcript of a BAG gene, or fragment thereof.

Detection of the DNA encoding BAG may also be used as an indicator of BAG expression. A plurality of
15 changes from wild type in the portion of DNA which constitutes a BAG gene may influence levels of gene expression. For example, gene amplification will provide more copies of a BAG gene within each cell, thereby facilitating the manufacture of an increased number of
20 mRNA molecules encoding BAG, which may result in an increased level of BAG protein within the cell. In another example, gene translocation or partial gene deletion may have an effect on the rate of gene expression by, for example, decreasing the ability of a
25 repressor protein to repress BAG transcription, resulting in higher levels of BAG protein within the cell. Detection of the portion of DNA which constitutes a BAG gene may have application by indicating the occurrence of gene amplification, translocation, mutation, partial
30 deletion, or other modification of the copy number, sequence, location, or accessibility of a BAG gene which results in abnormal levels of expression of a BAG gene.

Detection of the DNA constituting a BAG gene may take place by a variety of methods known to those of skill in the art. Such methods will typically include the detection of hybridization to the DNA constituting a BAG gene, or amplification of the DNA constituting a BAG gene. This method may be carried out by analysis of cellular DNA either *in vitro* or *in situ*. Numerous methods for carrying out this detection are known to those of ordinary skill in the art, as exemplified in the Current Protocols in Molecular Biology, and in U.S. Patent No. 5,882,864, which is incorporated herein in its entirety.

Classifying the patient

The classification of particular patients requires comparing the level of BAG gene expression in the tumor cells of a patient to a reference level (also referred to as a basal level). For example, after measuring the BAG expression level of the cells in a cancerous tumor of a patient, the measured level is compared to a reference level. This reference level is a level of expression of BAG used to evaluate the level of expression of BAG in cancerous cells of a patient. Specifically, when the level of BAG expression in the cancerous cells of a patient are higher than the reference level, the cells will be considered to have a high level of expression, or overproduction, of BAG. Conversely, when the level of BAG expression in the cancerous cells of a patient are lower than the reference level, the cells will be considered to have a low level of expression, or underproduction, of BAG.

As used herein, the terms "high level" or "overproduction" of BAG gene expression is related to a

level of BAG gene expression above a determined basal level and is likely different for each cancer type. Thus, in accordance with the present invention a reference or basal level of BAG gene expression in a particular cancer cell type is identified as a "cutoff" value, above which there is a significant correlation between the presence of BAG gene expression and increased or decreased tumor recurrence or spread. Those of skill in the art will recognize that some "cutoff" values are not sharp in that clinical correlations are still significant over a range of values on either side of the cutoff; however, it is possible to select an optimal cutoff value (e.g. varying H-scores, and the like) of BAG gene expression for each cancer cell type. It is understood that improvements in optimal cutoff values could be determined, depending on the sophistication of statistical methods used and on the number and source of samples used to determine reference or basal values for the different cancer cell types.

Such "overproduction" is not typically calculated in terms of absolute BAG gene expression or protein levels, but is determined using relative measurements. These relative measurements are illustrated for quantitation purposes with an "internal standard;" however, it will be appreciated that other standards or methods of determination may be used, such as comparison with external standards, BAG mRNA measurements, measurement of the DNA constituting a BAG gene, absolute values of the protein or mRNA or DNA levels, and the like.

The reference level may be determined by a plurality of methods, provided that the resulting reference level accurately provides a level of BAG

expression above which exists a first group of patients having a different risk of tumor recurrence or spread than that of a second group of patients having BAG expression levels below the reference level. The
5 reference level may be determined by, for example, measuring level of expression of BAG in non-tumorous cancer cells from the same tissue as the tissue of the cancer cells to be tested. The reference level may also be a level of BAG expression of *in vitro* cultured cells
10 which may or may not have been manipulated to simulate tumor cells, or may have been manipulated in any other manner which yields expression levels which accurately determine the reference level.

It is important to recognize that reference
15 levels are not necessarily the levels of BAG expression found in cultured cell lines used to provide internal standards. Rather, these may be reference level amounts of BAG gene expression (e.g., BAG protein, BAG mRNA, and the like) that occur in tumor cells. It is, for example,
20 possible that a high level of BAG gene expression in a tumor sample will have a relatively lower expression level than the corresponding BAG protein expression level in an internal standard.

The reference level may also be determined by
25 comparison of BAG expression levels in populations of patients having the same cancer. This may be accomplished by histogram analysis, in which the entire cohort of patients tested are graphically presented, wherein a first axis represents the level of BAG
30 expression, and a second axis represents the number of patients in the cohort whose tumor cells express BAG at a given level. Two or more separate groups of patients may be determined by identification of subsets populations of

the cohort which have the same or similar expression levels of BAG. Determination of the reference level may then be made based on an expression level which best distinguishes these separate groups.

5 Verification that the reference level distinguishes the likelihood of tumor recurrence or spread in cancer patients expressing below-reference levels of BAG versus cancer patients expressing above-reference levels of BAG is carried out using single
10 variable or multi-variable analysis. These methods determine the likelihood of a correlation between one or more variables and a given outcome. In the specific case, the methods will determine the likelihood of a correlation between BAG expression levels (or BAG
15 expression levels coupled with another variable) and disease-free or overall survival of cancer patients. Any one of a plurality of methods well known to those of ordinary skill in the art for carrying out these analyses may be used. Examples of single variable analysis is the
20 Kaplan-Meir method or the log-rank test. An example of multi-variable analysis is the Cox proportional-hazards regression model.

Population-based determination of reference levels (e.g., histogram analysis) is carried out using a
25 cohort of patients sufficient in size in order to determine two or more separate groups of patients having different BAG expression levels. Typically, such a cohort comprises at least 25 patients, preferably 50 patients, more preferably 75 patients, and yet more
30 preferably at least 100 patients. Similarly, verification of determined reference levels will also comprise at least 25 patients, preferably 50 patients,

more preferably 75 patients, and yet more preferably at least 100 patients.

The reference level may be a single number, equally applicable to every patient, or the reference level may vary, according to specific subpopulations of patients. For example, men might have a different reference level than women for the same cancer. Furthermore, the reference level might be some level determined for each patient individually. For example, the reference level might be a certain ratio of BAG expression in the tumor cells of a patient relative to BAG expression in non-tumor cells within the same patient. Thus the reference level for each patient may be proscribed by a reference ratio of BAG expression, wherein the reference ratio may be determined by any of the methods for determining the reference levels described above.

Further, while the reference level described above discusses a number which separates two groups of patients, it is within the scope of the invention that numerous reference values might exist which separate a plurality of populations. For example, two reference values may separate a first group of patients with high levels of BAG expression from a second group of patients with intermediate levels of BAG expression, and from a third group of patients with low levels of BAG expression. The number of different reference levels may be sufficient to proscribe a curve, such as a continuous line, which describes the likelihood of disease-free or overall survival in a patient as a function of BAG expression level in that patient. Such a curve will constitute a "continuous" BAG reference level, where the likelihood of disease free or overall survival in a

patient is portionional to the level of BAG expression level in that patient.

The reference level can also represent the level of BAG protein in one or more compartments of the cell. Typically, the reference level will represent the level of BAG protein in (a) the whole cell, (b) the nucleus, or (c) the cytosol. This level will be useful when cell compartmentalization of BAG protein correlates with the risk of tumor recurrence or spread of a certain cancer. For example, it has been observed that the level of BAG protein is higher in the cytosol of patients with low risk of breast cancer recurrence or spread than the cytosolic BAG protein level of patients with high risk of breast cancer recurrence or spread, despite the fact that the level of BAG expression in the nucleus can be roughly equivalent. Similarly, the reference level may be a ratio of levels of BAG protein in the different compartments (e.g., the ratio of nuclear BAG protein to whole cell BAG protein, or the ratio of nuclear BAG protein to cytosolic BAG protein).

It is also contemplated herein that the reference level can represent the level of expression of a single form of BAG (e.g., BAG-1N, BAG-1M, BAG-1L, BAG-2, BAG-3, BAG-4, OR BAG-5). This reference level will be useful when a single translated form of BAG correlates with the risk of tumor recurrence or spread of a certain cancer. For example, it may be observed that the level of BAG-1L expression is higher in patients with low risk of lung cancer recurrence or spread than the level of BAG-1L expression in patients with high risk of lung cancer recurrence or spread, despite the fact that the level of total BAG-1 expression (i.e., the sum of BAG-1N, BAG-1M, or BAG-1L expression levels) is the same

for all patients. Similarly, the reference level may be a ratio of levels of expression for different forms of BAG (e.g., the ratio of BAG-1N to BAG-1L, the ratio of BAG-2 to BAG-3, and the like).

5 The reference level of BAG expression may further be used in conjunction with another variable found to be a statistically significant indicator of the likelihood of disease-free or overall survival for a specific cancer. Such indicators include the presence or
10 levels of biomarkers of such as other proteins (e.g., ras, Bcl-2, and the like), or may be clinical or pathological indicators (e.g., age, tumor size, tumor histology, clinical stage, and the like). For example, clinical stage of the cancer is also be a statistically
15 significant indicator of disease-free or overall survival (in addition to BAG expression level), wherein the reference level of BAG expression may vary according to the clinical stage of the cancer. For example, levels of BAG expression, preferably high levels of BAG-1, in
20 conjunction with clinical stage I of a cancer for a given patient, are particularly preferred indicators for increased likelihood of disease free or overall survival.

Hence, the reference level of BAG expression may vary as a function of another statistically significant indicator
25 of disease-free or overall survival for a specific cancer.

Those of skill in the art will recognize that it is also possible to measure levels of BAG proteins in body fluid, such as serum. Tumors are known to readily
30 shed cells which, after release into the bloodstream, may burst due to cell fragility. For example, it is contemplated herein that overexpression of BAG-1 observed in approximately 2/3 of breast cancer patients results in

increased secretion and/or release of BAG-1 protein into serum. Thus, detection of any BAG levels, preferably high BAG-1 levels, present in body fluid (e.g., serum) is contemplated for use in the invention methods to
5 determine a prognosis of disease free or overall survival in a manner analogous to that demonstrated with the tissue samples. Very small quantities of BAG can be measured in body fluid, for example, using anti-BAG antibodies in immuno-PCR methods as described herein.

10 Accordingly, it is contemplated herein that the reference levels may represent the level of BAG present in a body fluid sample, such as serum. Invention methods that measure the level of circulating BAG (i.e., the level of BAG in blood or serum), preferably BAG-1, will
15 have a particularly preferred application to early diagnosis and screening, and early determination of risk of cancer recurrence or spread, for patients with abnormal levels of BAG (e.g., high levels) in their serum.

20 After the levels of expression of BAG in the tumor cells of a patient have been determined and compared to a reference level, the patient is then classified into a group having a certain likelihood of disease free or overall survival. Then the likelihood of
25 disease-free or overall survival for the patient is assessed based on the likelihood of disease-free or overall survival for patients in that group. For example, the tumor cells of a specific breast cancer patient may be determined to have high levels of BAG
30 expression relative to a reference level. This patient would then be classified into a group of patients having high levels of BAG expression. Since, in accordance with the present invention, it has been discovered that there

is an increased likelihood of disease-free or overall survival for the group of patients expressing high levels of BAG-1 in breast cancer cells (relative to those expressing low levels of BAG-1 in breast cancer cells),
5 the specific breast cancer patient would be considered to have an increased likelihood of disease free or overall survival.

In another embodiment of the present invention, a method is provided for prognosis of disease-free or
10 overall survival of individuals having a cancer tumor. In particular, the method comprises determining whether BAG protein is overproduced in a sample of such a tumor, wherein such an overproduction correlates positively with disease-free or overall survival. Thus, patients who
15 produce high levels of BAG protein have an increased likelihood of survival relative to patients producing low levels of BAG protein.

Another embodiment of the invention provides a method for screening a cancer patient to determine the
20 risk of tumor metastasis. The method comprises determining the level of amplification or expression of a BAG gene in a cancerous tissue sample from the patient. A patient found to have high levels of amplification or expression of the BAG gene, relative to a reference
25 level, is classified as being less likely to suffer tumor metastasis or having a increased chance of survival.

As used herein, the term "metastasis" refers to the growth of a cancerous tumor in an organ or body part, which is not directly connected to the organ of the
30 original cancerous tumor. Metastasis will be understood to include micrometastasis, which is the presence of an undetectable amount of cancerous cells in an organ or

body part which is not directly connected to the organ of the original cancerous tumor. Therefore, the present invention contemplates a method of determining the risk of further growth of one or more cancerous tumors in an organ or body part which is not directly connected to the organ of the original cancerous tumor.

In yet another embodiment of the invention, methods are provided for the determination of levels of BAG expression in cancer patients at an early stage of tumor development. As used herein, the term "stage", when applied to tumor development, refers to the degree of progression of a tumor. Various stages of tumor development are well known to those of skill in the art, as exemplified in Markman 1997, Basic Cancer Medicine. Stages of different cancers are defined according to different criteria, typically using the Tumor-Node-Metastasis (TNM) system. For example, stage I of breast cancer requires no lymph node involvement, and stage II of breast cancer involves either no lymph node involvement and a large primary tumor, or initial lymph node involvement and a small primary tumor (see Markman, supra, pages 35 and 36). Similar descriptions of the various clinical stages can be found in Markman, supra, for lung cancer (pages 54 to 55), and prostate cancer (pages 64 to 65), colon cancer (pages 78 to 79) and ovarian cancer (page 38 to 39). Early stages of tumor development shall be understood to refer to stages in tumor development in which the tumor has detectably spread no further than the lymph nodes local to the organ of the primary tumor. Typically, early stages will be considered to be stages I and II. As used herein, the phrase, "prior to lymph node involvement" refers to the detectable presence of cancer cells in the organ of the primary tumor, but the lack of a detectable presence of

cancer cells in any lymph nodes, including the lymph nodes closest to the organ of the primary tumor.

The predictive value of the method of the invention will be particularly effective in the case of patients in the early stages of cancer. This is because the method of the invention is advantageously effective in determining the risk of metastasis in patients who demonstrate no measurable metastasis at the time of examination. One of ordinary skill in the art would appreciate that the prognostic indicators of survival for cancer patients suffering from stage I cancer may be different from those for cancer patients suffering from stage IV cancer. For example, prognosis for stage I cancer patients may be oriented toward the likelihood of continued growth and/or metastasis of the cancer, whereas prognosis for stage IV cancer patients may be oriented toward the likely effectiveness of therapeutic methods for treating the cancer.

In another embodiment of the invention, a kit is provided to determine the levels of BAG expression in the cancerous tumor cells of the patient. Such a kit will comprise a reagent for detecting either the DNA encoding BAG, the mRNA encoding BAG, the BAG polypeptide, or any combination thereof. The reagent will comprise one or more molecules capable of specifically binding a nucleic acid sequence (DNA or RNA) encoding BAG, or the BAG polypeptide.

The kit may comprise one or more nucleic acid reagents for the detection of either DNA encoding BAG, mRNA encoding BAG, or both. The one or more nucleic acid reagents may be used for hybridization or amplification with the DNA and/or mRNA encoding BAG. The kit may

comprise one or more pairs of primers for amplifying the DNA and/or mRNA encoding BAG. The kit may further comprise samples of total mRNA derived from tissue of various physiological states, such as normal, and
5 metastatically progressive tumor, for example, to be used as controls. The kit may also comprise buffers, nucleotide bases, and other compositions to be used in hybridization and/or amplification reactions. Each solution or composition may be contained in a vial or
10 bottle and all vials held in close confinement in a box for commercial sale. Another embodiment of the present invention encompasses a kit for use in detecting the DNA and/or mRNA encoding BAG in cancer cells in a biological sample comprising oligonucleotide probes effective to
15 bind with high affinity to DNA and/or mRNA encoding BAG *in vitro* or *in situ* and containers for each of these probes.

In a further embodiment, the invention encompasses a kit for use in determining the level of BAG
20 expression in a biological sample comprising one or more agents, such as, for example, one or more antibodies, specific for one or more BAG polypeptides. In one particular embodiment, the kit will comprise one or more agents and one or more nucleic acid markers wherein the
25 agents and nucleic acid markers are modified in a fashion appropriate for carrying out immuno-polymerase chain reaction assays.

Another embodiment of the invention provides a method for determining the proper course of treatment for
30 a patient suffering from cancer. This method comprises determining the level of expression of a BAG gene in cancerous tissues from a patient. Then, a first group of patients is identified as having low levels of expression

of a BAG gene in the cells. This first group of patients may require treatment proper for patients having a lesser chance of survival or decreased time to tumor recurrence or spread. The method further comprises identifying a
5 second group of patients as having high levels of expression of a BAG gene in the cells. This second group of patients may require treatment proper for patients having a greater chance of survival and being less likely to suffer tumor recurrence or spread.

10 Appropriate treatments for patients having greater or lesser chances of survival will be determined, with proper consideration of other factors, by one of ordinary skill in the art according to the available therapeutic methods. As used herein, "course of
15 treatment" shall be understood to represent the treatment (e.g., surgery, chemotherapy, radiation, and the like) of a patient after diagnosis for cancer, and the span of time during which the patient is treated. Hence, a point in time considered to be "during the course of treatment"
20 shall represent any time between the initiation of the course of treatment and the termination of the course of treatment.

In a further embodiment of the invention, a method is provided for monitoring the effectiveness of a
25 course of treatment for a patient suffering from cancer. This method comprises:

- (a) determining a first BAG gene expression level in a cancerous tissue sample or a body fluid sample from said patient prior
30 to said treatment; and
- (b) subsequently determining a second BAG gene expression level in a cancerous tissue

sample or a body fluid sample from said patient during said treatment.

Comparison of said first BAG expression level with said second BAG expression level will then indicate the effectiveness of said treatment.

As used in the context of a course of treatment, "effectiveness" refers to the ability of the course of treatment to decrease the risk of tumor recurrence or spread and therefore to increase the likelihood of disease-free or overall survival of the patient. This method will have particular utility when the level of BAG expression in the tumor cells of a patient is abnormal compared to the level of BAG expression in the non-tumor cells of the patient.

Comparison of the first and second BAG expression levels will thereby serve to indicate whether BAG expression level is returning to that of non-tumor cells, implying a more effective course of treatment, or whether BAG expression level is remaining abnormal or increasing in abnormality, implying a less effective course of treatment. Levels of BAG expression may be determined using a plurality of samples from the patient, as described herein. A preferable sample form for this embodiment shall be a body fluid sample, such as serum sample, an exudate sample, and the like.

This embodiment of the invention is particularly useful when combined with the method of determining the risk of tumor recurrence or spread in a cancer patient, and thereby determining the proper course of treatment in a cancer patient. Specifically, the proper course of treatment of a cancer patient may be determined by determining the level of BAG expression in a sample from a patient, then classifying the patient's

likelihood of disease-free or overall survival according to the level of BAG expression. The course of treatment may then be monitored, according to the present embodiment of the invention, on one or more occasions to
5 determine the effectiveness of the course of treatment.

In a further embodiment of this invention, a method is provided for determining a prognosis of disease free or overall survival in a patient suffering from cancer. This method comprises:

- 10 (a) determining a BAG activity level in a cancerous tissue sample or body fluid sample from said patient; and
- (b) classifying said patient as belonging to either a first or second group of patients,
15 wherein said first group of patients having high levels of BAG activity is classified as having a different likelihood of suffering tumor recurrence or spread than said second group of patients having low levels of BAG
20 activity.

As used herein, "BAG activity level" or level of "BAG activity" refers to the level of active, uninhibited BAG polypeptides present in tumor cells or body fluid, and the degree of activity of these
25 polypeptides. Hence, level of activity shall be influenced by a plurality of factors including: levels of BAG, presence of less active or more active forms (including isoforms) of BAG, presence of less active or more active mutants of BAG, presence of proteins or other
30 molecules which increase BAG activity, presence of proteins or other molecules which decrease BAG activity. For example, levels of an antagonist to BAG, such as Hip (an antagonist to BAG-1, see for example Hohfeld and

Jentsch 1997, EMBO J 16:6029-6216), will be understood to commensurately lower BAG activity even if BAG protein levels remain unchanged. Determination of BAG activity shall therefore be carried out by a plurality of methods, including: assay of biological activity of BAG (such as ability to prevent cell death or other apoptotic assays, see, for example, United States Patent 5,550,019), assay of BAG protein levels, assay of the mRNA encoding a BAG protein, assay of the DNA which constitutes a BAG gene, assay of proteins (or the mRNA or DNA encoding the proteins) or other molecules which increase or decrease BAG activity, and any combination thereof.

All U.S. patents and all publications mentioned herein are incorporated in their entirety by reference thereto. The invention will now be described in greater detail by reference to the following non-limiting example.

EXAMPLES

This example illustrates the determination of the reference level of BAG-1 expression in breast cancer patients, and evaluation of the accuracy of the reference
5 level in predicting disease-free or overall survival of separate groups having different BAG-1 expression levels.

A retrospective correlative analysis was performed of BAG-1, Estrogen Receptor (ER), Progesterone Receptor (PR), HER2/Neu, p53, Bcl-2, Bax, which have all
10 been implicated in breast cancer formation. Immunohistochemical methods were carried out on archival paraffin blocks derived from 116 women with early-stage (stage I: n = 73, 63%; and stage-II: n = 43, 37%) breast cancer who were treated with lumpectomy followed by
15 radiation therapy to the intact breast using a median dose of 48 Grays (Gy) followed by an electron boost to the lumpectomy site to yield a total median dose of 64 Gy. There were 17/116 (15%) patients treated with adjuvant systemic chemotherapy and 20/116 (17%) patients
20 treated with tamoxifen therapy. The patients had a median follow-up of 12.4 years, with a minimum follow-up of 4 years (Table 1).

Table 1. Characteristics of breast cancer cases

DATA	No.	%
Number of patients	116	NA
Mean age (yrs.)	55	NA
5 Infiltrating ductal cancer	103	(89%)
Infiltrating lobular cancer	9	(8%)
Infiltrating medullary cancer	4	(3%)
Median follow-up (yrs.)	12.4	NA
10 Stage I/II	116	(100%)
Stage I	73	(63%)
Stage II	43	(37%)
Mean pathologic size (cm)	1.8	NA
Axillary dissection	62	(53%)
Positive lymph nodes	13	(11%)
15 Estrogen receptor positive	49	(42%)
Adjuvant chemotherapy	17	(15%)
Adjuvant tamoxifen	20	(17%)
Metastatic Disease	35	(30%)

20 The characteristics of patients used for this study are summarized: number (left column); percentage (right column). Metastatic disease indicates the number (and percentage) of patients who developed clinically detectable metastatic disease after diagnosis and treatment.

25 Immunostaining was carried out by colorimetric antibody detection using diaminobenzidine (brown) followed by hematoxylin (blue) counterstaining of nuclei. Immunostaining of the invasive component of breast cancers was scored according to intensity (0-4, wherein 0 is no staining and 4 is the most intense staining) and percentage of immunopositive cells (0-100%, for a minimum of 200 cells), evaluating the entire tissue-section, with the pathologist blinded to clinical details. Histo-

staining (H)-scores (0-400) were obtained by determining the product of intensity (0-4 scale) and percentage (0-100%). To set cut-offs for dichotomization of data into groups having high and low levels of expression, the H-score data for the entire data-set were displayed as dot-histograms with H-score on the x-axis and the number of patient samples having a given H-score on the y-axis. An H-score ≥ 150 was determined by this approach to be appropriate for use as a cut-off for high levels of BAG-1 expression. H-score data were displayed as histograms and dichotomized into high versus low expression level groups using optimized cut-offs for BAG-1 (H-score ≥ 150), Bcl-2 (H-score ≥ 180), and Bax (H-score ≥ 140). Immunoscoring for ER, PR, p53, and HER2/Neu was performed by established criteria (King et al., 1985, Cancer Research 45: 293-296).

BAG-1 is known to be present in the nuclei of some types of cells and nuclear immunostaining is normal for breast epithelium, whereas strong cytosolic immunostaining is abnormal here (Takayama et al. 1998, supra). This phenomenon was confirmed in the present example: compared to normal breast epithelium (NBE) cells, which were often present along with tumor cells in the same tissue-sections, cytosolic immunostaining for BAG-1 was clearly upregulated within the invasive breast cancer cells, with elevated levels of protein observed for 77/116 (66%) or roughly two-thirds of early-stage breast cancers compared with 9/83 (11%) of NBE ($p < 0.001$).

Some of the same tumor specimens also contained histologically evident ductal carcinomas in situ (DCIS); high levels of BAG-1 nuclear immunostaining were found in 9/12 (75%) and 6/12 (50%) sections had high levels of cytoplasmic BAG-1 protein levels in DCIS specimens,

suggesting that upregulation of BAG-1 can occur as a relatively early event in tumorigenesis and the translocation of protein from the nucleus to the cytoplasm may be important in cellular transformation.

5 Immunostaining data were correlated with distant metastasis-free survival (DMFS) and overall survival (OS). Kaplan-Meier analysis revealed that elevated levels of BAG-1 were significantly associated (i.e., $p \leq 0.05$) with longer DMFS ($p < 0.001$) and OS
10 ($p < 0.001$) (Figure 1). The group of patients having high BAG-1 protein levels (BAG-1 positive) experienced better 10-year DMFS than the group of patients having low (BAG-1 negative) BAG-1 protein levels (90% high group vs. 40% low group) and OS (84% high group vs. 40% low group).
15 Among the other biomarkers evaluated, only Bcl-2 was significant in univariate analysis as a predictor of longer DMFS ($p < 0.001$) and OS ($p < 0.001$). ER, PR, HER2/Neu, p53, and Bax were all insignificant in predicting survival for this cohort of patients. Among
20 clinical and pathological variables (e.g., age; tumor size; stage; tumor histology), only clinical stage I versus clinical stage II was significant in univariate analysis for predicting survival.

 In multivariate models using Cox-regression
25 analysis with variables including BAG-1, Bcl-2, Bax, p53, ER, PR, HER2/Neu, age and clinical stage, only BAG-1 retained statistical significance as a predictor of DMFS ($p = 0.008$) and OS ($p = 0.02$). All other biomarkers failed to reach clear statistical significance and
30 clinical stage (stage I versus stage II) was significant only for DMFS ($p = 0.029$) but not OS. Thus, patients whose tumors contain high levels of cytosolic BAG-1 protein are more likely to enjoy long-term survival and

freedom from tumor recurrence or spread and distant metastases, compared to those with tumors containing low levels of cytosolic BAG-1. These results demonstrate that BAG represents a novel and independent prognostic
5 factor which is associated with favorable outcome in patients with early-stage breast cancer.

Although the invention has been described with reference to the disclosed embodiments, it should be understood that various modifications can be made without
10 departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216